

# 

(43) International Publication Date 30 November 2000 (30.11.2000)

PCT

# (10) International Publication Number WO 00/71116 A1

(51) International Patent Classification<sup>7</sup>: C07D 207/34

A61K 31/40,

M., Dileep [IN/IN]; Vaishali Apartments, Pocket A-12, 3C, Kalkaji Extension, New Delhi 110 019, Maharashtra (IN).

- (21) International Application Number: PCT/IB00/00014
- (22) International Filing Date: 6 January 2000 (06.01.2000)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 775/DEL/99

25 May 1999 (25.05.1999) IN

- (71) Applicant (for all designated States except US): RAN-BAXY LABORATORIES LIMITED [IN/IN]; 19 Nehru Place, New Delhi 110 019, Maharashtra (IN).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): KUMAR, Yatendra [IN/IN]; U-26/5, DLF Qutab Enclave, Phase III, Gurgaon 122 001, Haryana (IN). THAPER, Rajesh, Kumar [IN/IN]; B-26, Sunshine Apartments, Block "C", Sushant Lok I, Gurgaon 122 002, Haryana (IN). KUMAR, S.,

- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
  - LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

0/71116 A1

#### (54) Title: PROCESS FOR THE PRODUCTION OF AMORPHOUS ATORVASTATIN CALCIUM

# PROCESS FOR THE PRODUCTION OF AMORPHOUS ATORVASTATIN CALCIUM

5

#### **FIELD OF THE INVENTION**

The present invention relates to a process for the production of amorphous atorvastatin calcium.

10

15

#### **BACKGROUND OF THE INVENTION**

Atorvastatin is chemically [R-(R\*,R\*)]-2-(4-fluoro-phenyl)-β dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H pyrrole-1-heptanoic acid. Atorvastatin calcium, a synthetic HMG-CoA reductase inhibitor, is used for the treatment of hyperlipidemia and hypercholesterolemia, both of which are risk factors for arteriosclerosis and coronary heart disease. Open dihydroxy carboxylic acid, lactone and various salt forms of atorvastatin have been synthesized.

United States Patent 5,273,995, describes that R-form of the ring opened acid form has surprising inhibition of the biosynthesis of cholesterol. Atorvastatin in its calcium salt form, i.e.  $[R-(R^*,R^*)]-2-(4-fluoro-phenyl)-\beta$ ,  $\delta$ -dihydroxy-5-(1-methylethyl)-3-phenyl-4- $[(phenylamino) \quad carbomyl]-1H-pyrrole-1-heptanoic acid calcium salt (2:1) having Formula 1:$ 

is more suited to formulations and has been recommended as a drug.

United States patents 5,003,080; 5,097,045; 5,103,024; 5,124,482; 5,149,837; 5,248,793; 5,280,126; 5,342,952, which are herein incorporated by reference, describe various processes and key intermediates for preparing atorvastatin.

Atorvastatin calcium produced by the processes described in the above mentioned United States patents does not give amorphous atorvastatin consistently but gives a mixture of its crystalline and amorphous forms, which has unsuitable filtration and drying characteristics and are not suitable for large-scale production.

20

5

10

15

PCT application, WO 97/03959, discloses novel crystalline forms of atorvastatin calcium designated as Form I, Form II, and Form IV and method for their preparation which provide more favourable filtration and drying characteristics.

PCT application WO 97/03960 describes a procedure for converting the crystalline form of atorvastatin to the amorphous form. Process disclosed therein comprises dissolving crystalline form-I atorvastatin in a non-hydroxylic solvent like tetrahydrofuran or mixtures of tetrahydrofuran and toluene. The process involves complete removal of the solvent under high temperature (about 90°C) and high vacuum (about 5mm) using capital intensive equipment. Exposure of the material to high temperature for several days leads to degradation of the product. This makes the process very inconvenient to operate at a large scale. Slow removal of solvents at a manufacturing scale renders this process as inefficient cost-wise and less productive.

10

5

#### **SUMMARY OF THE INVENTION**

It is an objective of the present invention to provide an efficient process for the production of amorphous atorvastatin, which eliminates the problems of prior art and is convenient to operate on a commercial scale.

15

Accordingly, the present invention provides a process for the preparation of atorvastatin calcium in an amorphous form which comprises dissolving crystalline atorvastatin in a non-hydroxylic solvent, adding a suitable non-polar hydro-carbon solvent and recovering atorvastatin from a solution thereof, by solvent precipitation, isolating and drying the product.

20

Generally, the product can be isolated by any standard method known in the art such as by filtration, centrifugation or decantation. Typically, this product is isolated by filtration when any of the solvents within the scope of the process are used.

Major advantages of the present invention compared to the prior art processes are:

- i. elimination of the need to remove solvent by drying techniques.
- ii. less time consuming with improved filtration.
- iii. easy to operate on large-scale.
- iv. reproducibly produces amorphous product having allowable levels of residual solvents.

The present invention thus provides a novel process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:

- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent to precipitate out the material; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium

The non-hydroxylic solvent is selected from a group of solvents, which have the ability to dissolve crystalline atorvastatin and includes tetrahydrofuran. Suitable non-polar hydrocarbon solvents are selected from a group consisting of: n-hexane, n-heptane, cyclohexane, hexane fraction, heptane fraction or the like. In a preferred embodiment of this invention, the non-hydroxylic solvent is tetrahydrofuran and antisolvent is n-hexane, cyclohexane or n-heptane.

Generally, crystalline atorvastatin calcium is dissolved in a non-hydroxylic solvent, e.g. tetrahydrofuran, at a concentration of about 15% w/v to about 40% w/v, preferably at a concentration of about 25% w/v to about 15% w/v at ambient

10

5

15

temperature and a non-polar hydrocarbon, preferably n-hexane, cyclohexane or n-heptane, is added at 0°C to 50°C, preferably at 20°C to 25°C. The product is recovered by filtration at ambient temperature. Filtration, which is fast and smooth, is carried out using nutsche filtration or centrifuge filtration. Preferably, nutsche filtration is used on large scale preparation. Filtered material, a semi-dry powder, is further dried to remove surface solvents in a vacuum tray drier, tray drier, fluid bed drier or a rotary vacuum drier to afford amorphous material. Preferably, material is dried in a vacuum tray drier at about 20°C to about 80°C for 6 hours to 24 hours. Most preferably, drying is carried out at about 50°C to about 60°C for 12 hours.

10

5

Quantity of antisolvent varies from 5 times to 50 times the input of crystalline atorvastatin calcium depending upon its solution in non-hydroxylic solvent. Preferably, the quantity of antisolvent used is about 20 times to about 40 times the input of crystalline atorvastatin calcium to make overall concentration of about 5% w/v to about 2.5 w/v%.

15

20

Amorphous atorvastatin calcium prepared according to the process of the present invention may be characterized by its x-ray powder diffration pattern (Figures 2) as shown in the accompanied drawings. X-ray powder diffration patterns (Figures 2) show no peaks which are characteristic of a crystalline atorvastatin calcium (Figure 1 of the accompanied drawings) thus demonstrating the amorphous nature of the product.

#### **BRIEF DESCRIPTION OF THE FIGURES**

Figure 1 is the diffractogram of crystalline atorvastatin calcium. The horizontal axis represents 2θ and the vertical axis corresponds to peak intensity.

Figure 2 is diffractogram of amorphous atorvastatin calcium. The horizontal axis represents 2θ and the vertical axis corresponds to peak intensity.

The present invention is illustrated by the following examples, which are not intended to limit the effective scope of the claims.

#### **DETAILED DESCRIPTION OF THE INVENTION**

#### Example 1

[R-(R\*,R\*)-2-(4-fluorophenyl)- $\beta$ ,  $\delta$ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium).

#### Method A

5

10

15

20

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. Clear solution so obtained was added slowly to cyclohexane (350 lt) under nitrogen atmosphere. It was vigorously stirred maintaining temperature at 20-25°C. The precipitated product was centrifuged and dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.5 kg) in an

amorphous form was obtained having residual solvent levels of 0.01% w/w tetrahydrofuran and 0.6% w/w cyclohexane. X-ray powder diffraction pattern (Figure 2 as shown in the accompanied drawings) demonstrate the amorphous nature of the product.

5

#### Method B

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. To a clear solution of atorvastatin, cyclohexane (350 lt) was added under vigorous stirring at 20 to 25°C. The precipitated mass was further stirred for 30 minutes and filtered in a centrifuge. The product was dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.6 kg) in an amorphous form was obtained having residual solvent levels of 0.01% w/w for tetrahydrofuran and 0.7% w/w for cyclohexane. X-ray powder diffraction pattern demonstrates the amorphous nature of the product.

15

10

#### Example 2

 $[R-(R^*,R^*)-2-(4-fluorophenyl)\beta, \qquad \delta-dihydroxy-5-(1-methylethyl)-3-phenyl-4-\\ [(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium)$ 

20

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-hexane instead of cyclohexane

to give amorphous atorvastatin (9.5 kg.). X-ray crystallography confirmed the amorphous nature of the product.

#### Example 3

 $[R-(R^*,R^*)]-2-4-fluorophenyl)-\beta, \qquad \delta-dihydroxy-5-(1-methylethyl)-3-phenyl-4-\\ [(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium)$ 

10

15

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-heptane instead of cylcohexane to give amorphous atorvastatin (9.6 kg). X-ray crystallography examination confirmed the amorphous nature of the product.


While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.


#### WE CLAIM:

 A process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:

- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent or adding the dissolved atorvastatin to the non-polar anti-solvent to precipitate out atorvastatin calcium; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium.
- 2. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is chosen from a group of non-polar hydrocarbon solvents comprising n-hexane, cyclohexane or n-heptane.
- 3. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-hexane.
- 4. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is cylcohexane.
- 5. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-heptane.
- 6. The process of claim 1, wherein said amorphous atorvastatin calcium is isolated by filtration.







Interr nal Application No PCT/IB 00/00014

		PCI/IB	00/00014
a classif IPC 7	REATION OF SUBJECT MATTER A61K31/40 C07D207/34		
According to	International Patent Classification (IPC) or to both national class	sification and IPC	
	SEARCHED		
Minimum do IPC 7	cumentation searched (classification system followed by classification sys	cation symbols)	
Ocumentati	ion searched other than minimum documentation to the extent t	hat such documents are included in the field	de searched
Electronic de	ata base consulted during the international search (name of dat	a base and, where practical, search terms	used)
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT	,	
Category *	Citation of document, with indication, where appropriate, of the	e relevant passages	Relevant to claim No.
A	WO 97 03959 A (WARNER LAMBERT CHRISTOPHER A (US); JENNINGS R 6 February 1997 (1997-02-06) cited in the application		1-6
A	DE 33 27 449 A (GLAXO GROUP LT 2 February 1984 (1984-02-02) page 9, line 7 - line 16 page 12, line 10 -page 13, lin		1-6
A	WO 97 03960 A (WARNER LAMBERT (US); SCHWEISS DIETER (US)) 6 February 1997 (1997-02-06) cited in the application claim 1	CO ;LIN MIN	1-6
		-/	
X Fur	ther documents are listed in the continuation of box C.	Patent family members are	listed in annex.
"A" docum	ategories of cited documents :  nent defining the general state of the art which is not idened to be of particular relevance	"T" later document published after the or priority date and not in conflic ched to understand the principle	t with the application but
"E" earlier filing "L" docum	document but published on or after the international date lent which may throw doubts on priority claim(s) or	invention "X" document of particular relevance cannot be considered novel or o involve an inventive step when t	annot be considered to
which citation "O" docum	n is cited to establish the publication date of another on or other special reason (as specified) nent referring to an oral disclosure, use, exhibition or r means	"Y" document of particular relevance carnot be considered to involve document is combined with one ments, such combination being	; the claimed invention an inventive step when the or more other such docu-
*P* docum	nent published prior to the international filing date but than the priority date claimed	in the art. "&" document member of the same p	·
Date of the	e actual completion of the international search	Date of mailing of the internation	nal search report
	10 May 2000	17/05/2000	
Name and	i mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL – 2280 HV Rijswijk	Authorized officer	
1	Tel. (+31-70) 340-2040, Tx. 31 651 epo nt, Fax: (+31-70) 340-3016	De Jong, B	

Intern tal Application No PCT/IB 00/00014

		rc1/18 00/00014
	RITION) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 03958 A (WARNER LAMBERT CO ; MCKENZIE ANN T (US)) 6 February 1997 (1997-02-06) cited in the application	1-6

information on patent family members

Inten nal Application No
PCT/IB 00/00014

	PCT/IB 00/00014			00/00014
Patent document cited in search report	Publication date	Patent family member(s)	<i>'</i>	Publication date
WO 9703959 A	06-02-1997	BR 96098 CA 22200 CN 11909 CZ 9800	296 A 187 A 372 A 318 A 955 A 121 A 705 A	18-02-1997 30-10-1998 23-03-1999 06-02-1997 19-08-1998 14-10-1998 24-06-1998
		HU 9900 IL 122 JP 11509 NO 980	207 A	30-04-1998 28-07-1999 14-07-1999 17-08-1999 16-01-1998
		SK 6	496 A 298 A 156 A	25-05-1998 07-10-1998 19-10-1999
DE 3327449 A	02-02-1984	AT 276 AU 566	154 B 783 A 881 B	26-01-1987 15-06-1986 05-11-1987
		BE 897 CA 1240 CH 657	783 A 422 A 313 A 134 A 528 B	02-02-1984 30-01-1984 09-08-1988 15-08-1986 14-02-1996
		CS 8305 CY 1 DE 3374	687 A 434 A 010 D 392 A	15-03-1988 02-09-1988 12-11-1987 25-05-1992
		DK 349 EP 0107 ES 524	083 A,B, 276 A 590 D	31-01-1984 02-05-1984 01-06-1985
		FI 832 FR 2531 GB 2127	689 A 757 A,B, 087 A 401 A,B	01-10-1985 31-01-1984 03-02-1984 11-04-1984
		HK 84 HU 190 IE 55	349 A 288 A 603 B 748 B	22-10-1984 28-10-1988 29-09-1986 02-01-1991
		IT 1168 JP 2025	375 A 3206 B 3666 C 3084 B	31-12-1986 20-05-1987 26-02-1996 05-04-1995
		KE 3 KR 9100	1391 A 18805 A 1046 B 1935 A	12-03-1984 03-06-1988 19-01-1991 23-11-1983
		MY 5 NL 8302 NO 832	5887 A 2705 A 2773 A,B, 5083 A	31-12-1987 16-02-1984
		PL 24: PT 7: SE 45:	3228 A 7135 A,B 3195 B	27-08-1984 01-08-1983 18-01-1988
		SG 26 SI 831	1208 A 5088 G 1558 A 3191 A	31-01-1984 15-07-1988 31-12-1995 11-07-1995

information on patent family members

Inten nal Application No PCT/IB 00/00014

	<del></del>	<del></del>	
Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3327449	4	SU 1266471 A	23-10-1986
		US 4994567 A	19-02-1991
		US 5013833 A	07-05-1991
		US 4562181 A	31-12-1985
		US 4820833 A	11-04-1989
		YU 155883 A	30-04-1986
		ZA 8305579 A	26-09-1984
		ZW 17383 A	26-10-1983
WO 9703960	A 06-02-1997	AU 700794 B	14-01-1999
		AU 6497896 A	18-02-1997
		BG 102188 A	31-08-1998
		BR 9609714 A	23-02-1999
		CA 2220455 A	06-02-1997
		CN 1190956 A	19-08-1998
		CZ 9800122 A	16-12-1998
		EP 0839132 A	06-05-1998
		HR 960312 A	28-02-1998
		IL 122161 A	14-07-1999
		JP 11510486 T	14-09-1999
		NO 980209 A	16-01-1998
		PL 324463 A	25-05-1998
		SK 5898 A	05-08-1998
WO 9703958	A 06-02-1997	AU 6484196 A	18-02-1997
		BG 102186 A	30-10-1998
		BR 9610567 A	06-07-1999
		CA 2220458 A	06-02-1997
		CN 1190957 A	19-08-1998
		CZ 9800123 A	17-06-1998
		EP 0848704 A	24-06-1998
		HR 960313 A	30-04-1998
		HU 9901687 A	28-10-1999
		IL 122162 A	14-07-1999
		JP 11509229 T	17-08-1999
		NO 980208 A	16-01-1998
			AA AC 1AAA
		PL 324532 A SK 5998 A	08-06-1998 06-05-1998